Introduction to Inelastic Neutron Scattering

Bruce D Gaulin
McMaster University

Brockhouse Institute for Materials Research



- Neutrons: Properties and Cross Sections
- Excitations in solids
- Triple Axis and Chopper Techniques
- Practical concerns





$^{235}U + n$

$\rightarrow$

daughter nuclei +

2-3 n + gammas

neutrons:

no charge

s=1/2

massive: mc^2~1 GeV

How do we produce neutrons



Fission

- chain reaction
- continuous flow
- 1 neutron/fission



Spallation

- no chain reaction
- pulsed operation
- 30 neutrons/proton


Front-End Building
Klystron Building
Linac Tunnel
Ring
Target
Future Target Building
Central Helium Liquefaction Building
Radio-Frequency Facility
Support Buildings
Center for Nanophase Materials Sciences
Joint Institute for Neutron Sciences
Central Laboratory and Office Complex
ORNL User Facility


Target
Compressor/ Accumulator
Accelerator
Pulsed Neutron Beam (TOF)
≈ 20 A
Current
Time
≈ 20 mA
≈ 1 ms
≈ 1 μs

Neutron interactions with matter

McMaster University

- **Properties of the neutron**
  - Mass m_n =1.675 x 10^{-27} kg
  - Charge 0
  - Spin-1/2, magnetic moment $\mu_n = -1.913\ \mu_N$

- **Neutrons interact with…**
  - Nucleus
  - Crystal structure/excitations (eg. Phonons)
  - Unpaired electrons via dipole scattering
  - Magnetic structure and excitations



Nuclear scattering



Magnetic dipole scattering

Wavelength-energy relations

- **Neutron as a wave …**
  - Energy (E), velocity (v), wavenumber (k), wavelength (λ)

$$k = \frac{m_n \mathrm{v}}{\mathrm{h}} = \frac{2\pi}{\lambda}$$

$$E = \frac{\mathrm{h}^2 k^2}{2m_n} = \frac{\mathrm{h}^2}{2m_n}\left(\frac{2\pi}{\lambda}\right)^2 = \frac{81.81\,\mathrm{meV}\cdot\mathrm{Å}^2}{\lambda^2}$$

$$E = k_B T = \left(0.08617\,\mathrm{meV}\cdot\mathrm{K}^{-1}\right)T$$



λ ~ interatomic spacing → E ~ excitations in condensed matter

	Energy (meV)	Temperature (K)	Wavelength (Å)
Cold	0.1 – 10	1 – 120	4 – 30
Thermal	5 – 100	60 – 1000	1 – 4
Hot	100 – 500	1000 – 6000	0.4 – 1

The Basic Experiment:



Incident Beam:

- **monochromatic**
- **“white”**
- **“pink”**

Scattered Beam:

- **Resolve its energy**
- **Don’t resolve its energy**
- **Filter its energy**

Fermi's Golden Rule within the 1st Born Approximation

$$W = 2\pi/h \; |\langle f | V | i \rangle|^2 \; \rho(E_f)$$

$$\delta\sigma = W / \Phi = (m/2\pi h^2)^2 \; k_f / k_i \; |\langle f | V | i \rangle|^2 \; \delta\Omega$$



$$\delta^2\sigma / \delta\Omega \, \delta E_f = k_f/k_i \; \sigma_{coh}/4\pi \; N \; S_{coh}(\mathbf{Q}, \omega)$$

$$+ k_f/k_i \; \sigma_{incoh}/4\pi \; N \; S_{incoh}(\mathbf{Q}, \omega)$$

Nuclear correlation functions



Pair correlation function

$$G(\mathbf{r},t) = \frac{1}{N}\int \sum_{jj'} \delta\left(\mathbf{r}' - \mathbf{R}_{j'}(0)\right)\delta\left(\mathbf{r}' + \mathbf{r} - \mathbf{R}_{j}(t)\right)d\mathbf{r}'$$

Intermediate function

$$I(\mathbf{Q},t) = \int G(\mathbf{r},t)e^{i\mathbf{Q}\cdot\mathbf{r}}d\mathbf{r} = \frac{1}{N}\sum_{jj'} \exp\left(-i\mathbf{Q}\cdot\mathbf{R}_{j'}(0)\right)\exp\left(i\mathbf{Q}\cdot\mathbf{R}_{j}(t)\right)$$

Scattering function

$$S(\mathbf{Q},\omega) = \frac{1}{2\pi\hbar}\int I(\mathbf{Q},t)e^{-i\omega t}dt$$

Differential scattering cross-section

$$\frac{d^2\sigma}{d\Omega dE_f} = \frac{\sigma_{scat}}{4\pi}\frac{k_f}{k_i}NS(\mathbf{Q},\omega)$$

Nuclear (lattice) excitations

McMaster University

Neutron scattering measures simultaneously the wavevector and energy of
collective excitations → dispersion relation, $\omega(\mathbf{q})$
In addition, **local excitations** can of course be observed

- **Commonly studied excitations**
  - Phonons
  - Librations and vibrations in molecules
  - Diffusion
  - Collective modes in glasses and liquids
- **Excitations can tell us about**
  - Interatomic potentials & bonding
  - Phase transitions & critical phenomena (soft modes)
  - Fluid dynamics
  - Momentum distributions & superfluids (eg. He)
  - Interactions (eg. electron-phonon coupling)

Atomic diffusion

For long times compared to the collision time, atom diffuses



$$\left\langle r^2(t)\right\rangle \approx 6Dt$$

Auto-correlation function

$$G_s(r,t) = \left\{6\pi\left\langle r^2(t)\right\rangle\right\}^{-3/2} \exp\left(-\frac{r^2}{6\left\langle r^2(t)\right\rangle}\right)$$

$$S(Q,\omega) = \frac{1}{\pi \mathrm{h}} \exp\left(\frac{\mathrm{h}\omega}{2k_BT}\right) \frac{DQ^2}{\omega^2 + (DQ^2)^2}$$

Liquid Na



Cocking, *J. Phys. C* **2**, 2047 (1969)..

Molecular vibrations

- Large molecule, many normal modes
- Harmonic vibrations can determine interatomic potentials



C60 molecule



Prassides *et al., Nature* **354**, 462 (1991).

Mapping Momentum – Energy (Q-E) space

Origin of reciprocal space;

Remains fixed for any sample rotation



Bragg diffraction:

Elastic scattering : $| k_i | = | k_f |$

Constructive Interference

Q = Reciprocal Lattice Vector



Bragg diffraction:

Constructive Interference

Q = Reciprocal Lattice Vector



Elastic scattering : $|k_i| = |k_f|$

Elementary Excitations in Solids



- Lattice Vibrations (Phonons)
- Spin Fluctuations (Magnons)



Energy vs Momentum

- Forces which bind atoms together in solids

Phonons

- Normal modes in periodic crystal → wavevector

$$\mathbf{u}(l,t) = \frac{1}{\sqrt{NM}} \sum_{j\mathbf{q}} \boldsymbol{\varepsilon}_j(\mathbf{q}) \exp(i\mathbf{q}\cdot\mathbf{l}) \hat{B}(\mathbf{q}j,t)$$

- Energy of phonon depends on **q** and polarization



Longitudinal mode Transverse mode



Lynn, *et al.*, *Phys. Rev. B* **8**, 3493 (1973).



FCC structure



FCC Brillouin zone

Phonon intensities

Structure (polarization) factor

$$S_{1+}(\mathbf{Q},\omega)=\frac{1}{2NM}e^{-Q^2\langle u^2\rangle}\sum_{j\mathbf{q}}\frac{\left|\mathbf{Q}\cdot\boldsymbol{\varepsilon}_j(\mathbf{q})\right|^2}{\omega_j(\mathbf{q})}\left(1+n(\omega)\right)\delta(\mathbf{Q}-\mathbf{q}-\boldsymbol{\tau})\delta\left(\omega-\omega_j(\mathbf{q})\right)$$





Guthoff *et al., Phys. Rev. B* **47**, 2563 (1993).

More complicated structures



Woods, *et al., Phys. Rev*. **131**, 1025 (1963).



Chaplot, *et al., Phys. Rev*. B **52**, 7230(1995).

Acoustic phonon



Optical phonon



La_2CuO_4

Spin excitations

McMaster University

- **Spin excitations**
  - Spin waves in ordered magnets
  - Paramagnetic & quantum spin fluctuations
  - Crystal-field & spin-orbit excitations

- **Magnetic inelastic scattering can tell us about**
  - Exchange interactions
  - Single-ion and exchange anisotropy (determine Hamiltonian)
  - Phase transitions & critical phenomena
  - Quantum critical scaling of magnetic fluctuations
  - Other electronic energy scales (eg. CF & SO)
  - Interactions (eg. spin-phonon coupling)

Spin waves

Ferromagnetic



Perring *et al., Phys. Rev*. Lett. **77**, 711 (1996).

Antiferromagnetic



Shapiro *et al., Phys. Rev*. B **10**, 2014 (1974).

Ferrimagnetic



McQueeney *et al., Phys. Rev*. Lett. **99**, 246401 (2007).

Scattering experiments



Single-crystal



Powder S(|**Q**|,ω)



Instrument and sample (powder or single-crystal) determine how (**Q**,ω) space is sampled







Bragg's Law: $n\lambda = 2d \sin(\theta)$







Bragg's Law: $n\lambda = 2d \sin(\theta)$

Brockhouse's Triple Axis Spectrometer

$| k_i | = 2 \pi / \lambda_i$



$| k_f | = 2 \pi / \lambda_f$



Momentum Transfer:

$Q = k_i - k_f$





Energy Transfer:

$\delta E = h^2/2m\,(k_i^2 - k_f^2)$

Two Axis Spectrometer:

- 3-axis with analyser removed

- Powder diffractometer
- Small angle diffractometer
- Reflectometers

Diffractometers often employ working assumption that all scattering is *elastic.*





Soller Slits: Collimators

Define beam direction to +/- 0.5, 0.75 etc. degrees





Filters:
Remove λ /n from incident or scattered beam, or both







Single crystal monochromators:

Bragg reflection and harmonic contamination





$n\lambda = 2d \sin(\theta)$

Get: λ , $\lambda/2$, $\lambda/3$, etc.

Pyrolitic graphite filter:





E = 14.7 meV

λ = 2.37 A

v = 1.6 km/s

2 x v = 3.2 km/s

3 x v = 4.8 km/s



Constant k_f (ii)



Constant k_i

Two different ways of performing constant-Q scans

Mapping Momentum – Energy (Q-E) space

Origin of reciprocal space;

Remains fixed for any sample rotation




$-k_1$
k_o
Q
(i i)


$-k_1$
k_o
Q
(i,j)


Collimator
Reactor
Shielding drum
Monochromator crystal
Filter
Monitor
Soller slits
Analyzer crystal
Boral shielding
Soller slits
Shielding
Detector

Elementary Excitations in Solids



- Lattice Vibrations (Phonons)
- Spin Fluctuations (Magnons)



Energy vs Momentum

- Forces which bind atoms together in solids





Constant Q, Constant E
3-axis technique allow us to
Put Q-Energy space on a grid,
And scan through as we wish

Map out elementary excitations
In Q-energy space (dispersion
Surface)

Samples

McMaster University

- **Samples need to be BIG**
  - ~ gram or cc
  - Counting times are long (mins/pt)



- **Sample rotation**
- **Sample tilt**





Co-aligned $CaFe_2As_2$ crystals

Monochromators

McMaster University

- **Selects the incident wavevector**



$$Q(hkl) = \frac{2\pi}{d(hkl)} = 2k_i \sin\theta$$



- Reflectivity
- focusing
- high-order contamination
  eg. λ/2 PG(004)

Mono	d(hkl)	uses
PG(002)	3.353	General
Be(002)	1.790	High k_i
Si(111)	3.135	No λ/2

Detectors

- **Gas Detectors**
- n + ^{3}He → ^{3}H + p + 0.764 MeV
- Ionization of gas
- e^- drift to high voltage anode
- High efficiency



- **Beam monitors**
- Low efficiency detectors for measuring beam flux

Resolution

- **Resolution ellipsoid**
  - Beam divergences
  - Collimations/distances
  - Crystal mosaics/sizes/angles



- **Resolution convolutions**

$$\mathrm{I}(\mathbf{Q}_0,\omega_0) = \int \mathrm{S}(\mathbf{Q}_0,\omega_0) R(\mathbf{Q}-\mathbf{Q}_0,\omega-\omega_0)\, d\mathbf{Q}\, d\omega$$

Resolution focusing

- **Optimizing peak intensity**
- **Match slope of resolution to dispersion**



Neutrons have *mass*
so higher energy means faster – lower energy means slower



v (km/sec) = 3.96 / λ (A)

- 4 A neutrons move at ~ 1 km/sec
- DCS: 4 m from sample to detector
- It takes 4 msec for elastically scattered 4 A neutrons to travel 4 m

- msec timing of neutrons is easy
- δ E / E ~ 1-3 % - very good !

We can measure a neutron's energy, wavelength by measuring its *speed*

Time-of-flight methods

Spallation neutron source



Pharos – Lujan Center

- Effectively utilizes time structure of pulsed neutron groups

$$t = \frac{d}{v} = \left(\frac{m}{h} d\right)\lambda$$






Optical Filter
Crystal Filter
White Beam Monitor
Monochromatic Beam Monitor
Sample Chamber
Radial Collimator
Transmitted Beam Monitor
Sample
Neutron Guide
Shutter
Frame Overlap Chopper
Order Removal Choppers
Pulsing Choppers
Monochromating Choppers
Argon gas-filled Flight Chamber
Beam Stop
913 3He detectors, 400x31x11mm3

A single (disk) chopper pulses the neutron beam.



A second chopper selects neutrons within a narrow range of speeds.



Counter-rotating choppers (close together), with speed ◆, behave like single choppers with speed 2◆. They can also permit a choice of pulse widths.

Additional choppers remove "contaminant" wavelengths and reduce the pulse frequency at the sample position.

The DCS has seven choppers, 4 of which have 3 "slots"



Disk 4B

Fermi Choppers

- Body radius ~ 5 cm
- Curved absorbing slats
  - B or Gd coated
  - ~mm slit size
- f = 600 Hz (max)
- Acts like shutter, $\Delta t \sim \mu s$





Figure 1. ISIS MAPS chopper and slit package assembly – exploded view



T-zero chopper

- **Background suppression**
- **Blocks fast neutron flash**





Position sensitive detectors

- ^{3}He tubes (usu. 1 meter)
- Charge division
- Position resolution ~ cm
- Time resolution ~ 10 ns

MAPS detector bank



Sample environment

McMaster University

- **Temperature, field, pressure**
- **Heavy duty for large sample environment**
  - CCR
  - He cryostats
  - SC magnets
  - …

- Can be machined from Al

  ~ neutron transparent

  relatively easy to work with

Guides

- **Transport beam over long distances**
- **Background reduction**
- **Total external reflection**
  - Ni coated glass
  - Ni/Ti multilayers (supermirror)



Size matters



SEQUOIA detector vacuum vessel

- **Length = resolution**
  - Instruments ~ 20 – 40 m long
  - E-resolution ~ 2-4% E_i
- **More detectors**
  - SEQUOIA – 1600 tubes, 144000 pixels
  - Solid angle coverage 1.6 steradians
- **Huge data sets**
- 0.1 – 1 GB

Kinematic limitations

- **Many combinations of k_i,k_f for same Q,ω**
  - Only certain configurations are used (eg. E_f-fixed)

- **Cannot "close triangle" for certain Q,ω due to kinematics**



Minimum accessible Q



Kinematic limits, E_i=160 meV

Data visualization

- Large, complex data from spallation sources
- Measure S(**Q**,ω) – 4D function

$La_{1-x}Ca_xMnO_3$



Ye *et al.*, *Phys. Rev. B*, **75** 144408 (2007).





Development of Long Range Order



The appearance of spin waves indicates that the field-induced state is long range ordered

Field-induced order in the Pyrochlore Yb2Ti2O7:

Weak magnetic field // [110] induces LRO

appearance of long-lived spin waves
at low T and moderate H



References

General neutron scattering

G. Squires, "Intro to theory of thermal neutron scattering", Dover, 1978.

S. Lovesey, "Theory of neutron scattering from condensed matter", Oxford, 1984.

R. Pynn, http://www.mrl.ucsb.edu/~pynn/.

Polarized neutron scattering

Moon, Koehler, Riste, Phys. Rev **181**, 920 (1969).

Triple-axis techniques

Shirane, Shapiro, Tranquada, "Neutron scattering with a triple-axis spectrometer", Cambridge, 2002.

Time-of-flight techniques

B. Fultz, http://www.cacr.caltech.edu/projects/danse/ARCS_Book_16x.pdf